JLAB-THY-12-1644

arXiv:1210.2364v1 [nucl-th] 8 Oct 2012

Baryon Masses and Axial Couplings in the Combined $1/\mathbf{N_c}$ and Chiral Expansions

A. Calle Cordón[1,*] and J. L. Goity[1,2,†]

[1] *Thomas Jefferson National Accelerator Facility,*
Newport News, Virginia 23606, USA.
[2] *Department of Physics, Hampton University, Hampton, VA 23668, USA.*

Abstract

The effective theory for baryons with combined $1/N_c$ and chiral expansions is analyzed for non-strange baryons. Results for baryon masses and axial couplings are obtained in the small scale expansion, to be coined as the ξ-expansion, in which the $1/N_c$ and the low energy power countings are linked according to $1/N_c = \mathcal{O}(\xi) = \mathcal{O}(p)$. Masses and axial couplings are analyzed to $\mathcal{O}(\xi^3)$ and $\mathcal{O}(\xi^2)$ respectively, which correspond to next-to-next to leading order evaluations, and require one-loop contributions in the effective theory. The role of the spin-flavor approximate symmetry in baryons, consequence of the large N_c limit, is manifested in the physical world with $N_c = 3$ in a significant way, as the analysis of its breaking in the masses and the axial couplings shows. Applications to the recent lattice QCD results on baryon masses and the nucleon's axial coupling are presented. It is shown that those results are naturally described within the effective theory at the order considered in the ξ-expansion.

PACS numbers: 11.15-Pg, 11.30-Rd, 12.39-Fe, 14.20-Dh

Keywords: Baryons, large N, Chiral Perturbation Theory

---

[*] Electronic address: cordon@jlab.org

[†] Electronic address: goity@jlab.org

Contents

I. Introduction 2

II. Framework for the combined $1/\mathrm{N_c}$ expansion and Baryon Chiral Perturbation Theory 4

A. Consistency of the $1/N_c$ expansion 7

B. ξ power counting 8

III. Baryon masses 10

IV. Axial couplings 13

V. Analysis of lattice QCD results for baryon masses and the nucleon's axial coupling 16

VI. Discussion and conclusions 22

Acknowledgments 23

A. Spin-flavor Algebra 24

B. Non-linear realization of chiral symmetry and spin-flavor transformations 24

C. Tools for building effective Lagrangians 25

D. Matrix elements of spin-flavor operators in the symmetric representations of SU(4) 26

1. Useful matrix elements 27

References 29

I. INTRODUCTION

The low energy effective theory for baryons is a topic that has evolved over time through several approaches and improvements. The early version of baryon Chiral Perturbation Theory (ChPT) [1] evolved into the various effective field theories based on effective chiral

Lagrangians [2–4], starting with the relativistic version [5, 6] or Baryon ChPT (BChPT), followed by the non-relativistic version based in an expansion in the inverse baryon mass [6, 7] or Heavy Baryon ChPT (HBChPT), and by manifestly Lorentz covariant versions based on the IR regularization scheme [8–10]. In all these versions of the baryon effective theory a consistent low energy expansion can be implemented. The most important issue, which became apparent quite early, was the convergence of the low energy expansion. Being an expansion that progresses in steps of $\mathcal{O}(p)$ in contrast to the expansion in the pure Goldstone Boson sector where the steps are $\mathcal{O}(p^2)$, it is natural to expect a slower rate of convergence. However, a key factor with the convergence has to do with the important effects due to the closeness in mass of the spin 3/2 baryons. It was realized [11], that the inclusion of those degrees of freedom play an important role in improving the convergence of the one-loop contributions to certain observables such as the π-N scattering amplitude and the axial currents and magnetic moments. There have been since then numerous works including spin 3/2 baryons [12–20]. The key enlightenment resulted from the study of baryons in the large N_c limit of QCD [21]. It was realized that in that limit baryons behave very differently than mesons [22], in particular because their masses scale like $\mathcal{O}(N_c)$ and the π-baryon couplings are $\mathcal{O}(\sqrt{N_c})$. These properties were shown to require for consistency that at large N_c baryons must respect a dynamical contracted spin flavor symmetry $SU(2N_f)$, N_f being the number of light flavors [23–26], broken by effects ordered in powers of $1/N_c$ and in the quark mass differences. The inclusion of the consistency requirements of the large N_c limit into the effective theory came naturally through a combination of the $1/N_c$ expansion and HBChPT [27], which is the framework followed in the present work. The study of one-loop corrections in that framework was first carried out in Refs. [27–29]. In the combined theory one has to deal with the fact that the $1/N_c$ and Chiral expansions do not commute [30]. The reason is due to the presence of the baryon mass splitting scale of $\mathcal{O}(1/N_c)$ ($\Delta - N$ mass difference), for which it becomes necessary to specify its order in the low energy expansion. Thus the $1/N_c$ and Chiral expansions must be linked. Particular emphasis will be given to the specific linking in which the baryon mass splitting is taken to be $\mathcal{O}(p)$ in the Chiral expansion, and which will be called the ξ-expansion. Following references [27–29], the theoretical framework is presented here in detail, in particular the renormalization, the power countings, and the linked $1/N_c$ and low energy expansions, along with observations that further clarify the significance of the framework.

The very significant contemporary progress in the calculations of baryon observables in lattice QCD (LQCD) [31–33] opens new opportunities for further understanding the low energy effective theory of baryons. The determination of the quark mass dependence of the various low energy observables, such as masses, axial couplings, magnetic moments, electromagnetic polarizabilities, etc., are of key importance as a significant test of the effective theory, in particular its range of validity in quark masses, as well as for the determination of its low energy constants (LECs). Lattice results for the N and Δ masses [34–39] and the axial coupling g_A of the nucleon [40–45] at varying quark masses are analyzed with the purpose of testing the effective theory presented here. This in turn can give insights on LQCD results, in particular an understanding on the role and relevance of including the spin 3/2 baryons consistently with large N_c requirements.

This work is organized as follows. In Section II the framework for the combined $\mathbf{1}/\mathrm{N_c}$ and HBChPT expansions is presented. Section III presents the evaluation of the baryon masses and Section IV the one for axial couplings at the one-loop level. Section V is devoted to applying those results in the ξ-expansion to LQCD results. Finally, Section VI is devoted to observations and conclusions . Several appendices present useful material used in the calculations, namely, Appendix A on spin-flavor algebra, Appendix B on symmetries, Appendix C on the construction of effective Lagrangians, and Appendix D on useful matrix elements of spin-flavor operators.

II. FRAMEWORK FOR THE COMBINED $1/\mathrm{N_c}$ EXPANSION AND BARYON CHIRAL PERTURBATION THEORY

In this section the framework for the combined $1/N_c$ and chiral expansions in baryons is presented in some detail along similar lines as in the original works [27–29]. The symmetries that the effective theory must respect in the chiral and large N_c limits are chiral $SU_L(N_f) \times SU_R(N_f)$ and contracted dynamical spin-flavor symmetry $SU(2N_f)$[23–26] [1]. N_f is the number of light flavors, and in this work $N_f = 2$. In the limit $N_c \to \infty$ the spin-flavor symmetry requires baryons to belong into degenerate multiplets of $SU(4)$. In particular, the ground state (GS) baryons belong into a symmetric $SU(4)$ multiplet, which consists of states with $I = S$, being S the baryon spin and I its isospin. At finite N_c the spin-flavor symmetry is broken by effects suppressed by powers of $1/N_c$, and the baryon mass splittings in the GS multiplet are proportional to $(S+1)/N_c$. The effects of finite N_c are then implemented as an expansion in $1/N_c$ at the level of the effective Lagrangian. Because baryon masses scale as proportional to N_c, it becomes natural to use the framework of HBChPT [7, 46], where the expansion in inverse powers of the baryon mass becomes part of the $1/N_c$ expansion. The framework presented next follows that of Refs. [27, 28].

---

[1] See also Appendix B.

The non-relativistic baryon field, denoted by $\mathbf{B}$, consists of the symmetric spin-flavor $SU(4)$ multiplet with states $I = S$, $S = 1/2, \cdots, N_c/2$ (N_c odd). Chiral symmetry is realized in the usual non-linear way on $\mathbf{B}$, namely [2–4]:

$$(L, R) : \mathbf{B} = h(L, R, u)\mathbf{B}, \tag{1}$$

where $L(R)$ is a $SU_{L(R)}(2)$ transformation, u is given in terms of the pion fields π^a by $u = \exp(i\pi^a\tau^a/2F_\pi)$, $F_\pi = 92.4$ MeV, and $h(L, R, u)$ is an $SU_I(2)$ isospin transformation which in any representation of Isospin satisfies $R\,u\,h^\dagger(L, R, u) = h(L, R, u)\,u\,L^\dagger$. The chiral covariant derivative $D_\mu\mathbf{B}$ is given by:

$$\begin{aligned} D_\mu\mathbf{B} &= \partial_\mu\mathbf{B} - i\Gamma^a_\mu\hat{I}^a\mathbf{B}, \\ \Gamma_\mu &= \frac{1}{2}(u^\dagger(i\partial_\mu + r_\mu)u + u(i\partial_\mu + l_\mu)u^\dagger), \\ \Gamma^a_\mu &= \frac{1}{2}\langle\tau^a\Gamma_\mu\rangle, \end{aligned} \tag{2}$$

where $l_\mu = v_\mu - a_\mu$ and $r_\mu = v_\mu + a_\mu$ are gauge sources. Here the notation $\langle A\rangle \equiv \mathrm{Tr}A$ is used for flavor traces, and in general, $A^a = \frac{1}{2}\langle\tau^a A\rangle$ where A is in the fundamental representation in the trace. The definition $\tau^0 = I_{2\times2}$ is used. The axial Maurer-Cartan one-form necessary as building block of the effective chiral Lagrangian is:

$$u_\mu = u^\dagger(i\partial_\mu + r_\mu)u - u(i\partial_\mu + l_\mu)u^\dagger, \qquad (L, R) : u_\mu = h(L, R, u)u_\mu h^\dagger(L, R, u). \tag{3}$$

Since $F_\pi = \mathcal{O}(\sqrt{N_c})$, u, u_μ, Γ_μ contain different orders in the expansion in powers of $1/N_c$. The contracted $SU(4)$ transformations (see Appendix A) are generated by $\{S^i, I^a, X^{ia}\}$, where $X^{ia} = G^{ia}/N_c$ are semiclassical at large N_c. The ordering in N_c of the matrix elements of the spin-flavor generators in states with $S = \mathcal{O}(N_c^0)$ are as follows: $S^i = \mathcal{O}(N_c^0)$, $I^a = \mathcal{O}(N_c^0)$, and $G^{ia} = \mathcal{O}(N_c)$. While infinitesimal $SU(4)$ transformations generated by I^a

correspond to the usual isospin transformations when acting on pions, the ones generated by X^{ia} affect only the baryons (one can define these generators to not affect the pion field as shown in Appendix B). The effective Lagrangian can be systematically written as a power series in the low energy expansion or Chiral expansion, and simultaneously in $1/N_c$. It is most convenient to write the Lagrangian to be manifestly chiral invariant as is usually done. The low energy constants (LECs) will themselves admit an expansion in powers of $1/N_c$. For the HBChPT expansion the large mass of the expansion is taken to be the spin-flavor singlet component of the baryon masses, $M_0 = N_c\, m_0$ (m_0 can be considered here to be a LEC defined in the chiral limit and which will have itself an expansion in $1/N_c$). To $\mathcal{O}(1/N_c)$ baryon masses will read [25, 26]:

$$m_{\mathbf{B}}(S) = M_0 + \frac{C_{HF}}{N_c} S(S+1) + c_1\, N_c\, M_\pi^2 + \cdots . \tag{4}$$

The baryon mass splittings due to the hyperfine term, second term in Eq. (4), must be considered to be a small energy scale. It becomes necessary to establish of what order that term is in the low energy expansion, as it naturally appears in combinations with powers of M_π when loop diagrams are calculated. This fact makes that the low energy and $1/N_c$ expansions do not commute [30, 47], and the natural way to proceed is therefore to link the two expansions. For the purpose of organizing the effective Lagrangian it is convenient to established the link between the two expansions. In the real world with $N_c = 3$ the $\Delta - N$ mass splitting is about 300 MeV, and therefore it is reasonable to count that quantity as $\mathcal{O}(p)$ in the low energy expansion: the expansion where $1/N_c = \mathcal{O}(p) = \mathcal{O}(\xi)$ will be adopted in what follows, and it will be called ξ-expansion. This power counting corresponds to the so called small scale expansion (SSE) [13], now consistently implemented in the context of the $1/N_c$ expansion. Whenever appropriate, it will be indicated which aspects of the analysis are general and which are only valid in that expansion. Up to $\mathcal{O}(\xi)$ the baryon effective Lagrangian reads:

$$\mathcal{L}_{\mathbf{B}}^{(1)} \; = \; \mathbf{B}^\dagger \left(iD_0 + \mathring{g}_A u_{ia} G^{ia} - \frac{C_{HF}}{N_c} \hat{S}^2 - \frac{c_1}{2} N_c\, \chi_+ \right) \mathbf{B}, \tag{5}$$

where $\mathring{g}_A$ is the axial coupling in the chiral and large N_c limits (it has to be rescaled by a factor 5/6 to coincide with the usual axial coupling as defined for the nucleon), χ_+ is the source containing the quark masses: specifically $\chi_+ = 2M_\pi^2 + \cdots$ (see Appendix C). Here one notes an important point which will be present in other instances as well: the baryon mass dependence on the current quark mass behaves at $\mathcal{O}(N_c\, M_\pi^2)$ (c_1 is of zeroth order in N_c), and this indicates that in a strict large N_c limit the expansion in the quark masses of certain quantities such as the baryon masses cannot be defined due to divergent coefficients of $\mathcal{O}(N_c)$. The Lagrangian is manifestly invariant under the chiral transformations, translations and rotations (the latter also involving obviously the action of the S^i generators of $SU(4)$). Under transformations generated by X^{ia}, the kinetic term changes at $\mathcal{O}(1/N_c^2)$, the term proportional to $\mathring{g}_A$ which contains the $\pi\mathbf{B}\mathbf{B}'$ interaction and the leading order terms of the axial currents, changes by terms which are a factor $\mathcal{O}(1/N_c^2)$ smaller than the original term, the term proportional to c_1, which gives the leading order (LO) σ-term in the baryon masses, is a spin-flavor singlet, and finally the hyperfine term proportional to C_{HF} is the only one that shows explicit spin-flavor symmetry breaking as it changes by $\mathcal{O}(1/N_c)$, which is the order of the term itself (this is so because $[\hat{S}^2, X^{ia}] = \mathcal{O}(N_c^0)$). That term therefore provides the dominant spin-flavor symmetry breaking in the effective Lagrangian. The higher order Lagrangians can be built using the tools provided in Appendix C . The operators appearing in the effective Lagrangian are normalized in such a way that all the LECs are of zeroth order in N_c. The $1/N_c$ power ν_{1/N_c} of the operators in the Lagrangian are determined by the following simple rule:

$$\nu_{1/N_c} = n - 1 - \kappa + \frac{n_\pi}{2}, \tag{6}$$

where the spin-flavor operator is n-body (n is the number of factors of $SU(4)$ generators appearing in the operator), κ is basically the number of factors of the generators G^{ia} remaining after reducing the operator using commutators, and n_π is the number of pions attached to the vertex. It is opportune to point out that commutators of spin-flavor operators will always reduce the n-bodyness of the product of operators: e.g., let $\mathcal{G}$ be any generator of $SU(4)$, and consider the commutator $[\mathcal{G}, \hat{S}^2] = \{S^i, [\mathcal{G}, S^i]\}$, and then, because $[\mathcal{G}, S^i]$ is a 1-body operator, $[\mathcal{G}, \hat{S}^2]$ is actually a 2-body operator.

A. Consistency of the $1/N_c$ expansion

The consistency of the $1/N_c$ expansion in QCD gives rise to the dynamical spin-flavor contracted $SU(2N_f)$ symmetry in baryons at large N_c. At the baryon level that symmetry can be deduced as the result of consistency or correct N_c power counting, of observables in which pion-baryon couplings are involved: because the pion-baryon coupling is $\mathcal{O}(\sqrt{N_c})$

from Witten's counting rules [22]. In particular the consistency of pion-baryon scattering is a direct way of deriving the existence of the dynamical spin-flavor symmetry [25, 26]. In general, for any quantity there must be cancellations between the terms with the "wrong" power counting stemming from different Feynman diagrams. For instance, baryon masses are $\mathcal{O}(N_c)$, and therefore pion loop contributions cannot give contributions which scale with a higher power of N_c. On the other hand, the baryon mass splittings are $\mathcal{O}(1/N_c)$, and loop contributions must respect that scaling. Similarly, in the axial currents, whose matrix elements are $\mathcal{O}(N_c)$ such cancellations occur when loop corrections are calculated. All this will be illustrated in the application to baryon masses and axial couplings discussed next. Although certain key cancellations must be exact in the large N_c limit, the analysis of LQCD results will show that they are very significant in the physical world where $N_c = 3$.

B. ξ power counting

The terms in the effective Lagrangian are constrained in their N_c dependence by the requirement of the consistency of QCD at large N_c. This constraint is in the form of a lower bound in the power in $1/N_c$ for each term one could write down in the Lagrangian. This leads to constraints on the N_c dependencies of the ultra-violet (UV) divergencies, which have to be subtracted by the corresponding counter-terms in the Lagrangian. One very important point to mention is that the UV divergencies are necessarily polynomials in low momenta p (derivatives), in M_π^2 and in $1/N_c$ (modulo factors of $1/\sqrt{N_c}$ due to $1/F_\pi$ factors in terms where pions are attached). Therefore, the structure of counter-terms is independent of any linking between the $1/N_c$ and chiral expansions. For this reason, one can simply take the large N_c and low energy limits independently in order to determine the UV divergencies. The Chiral or low energy order of a diagram is given by [48]:

$$\nu_p = 2 - n_B/2 + 2L + \sum_i n_i(\nu_{p_i} + b_i/2 - 2), \tag{7}$$

where n_B is the total number of external baryon legs (two in the present work), L the number of loops, n_i the number of vertices of type i, ν_{p_i} is the the low energy (Chiral) power of that type of vertex, and b_i the number of baryon legs of the vertex (0 or 2 in the single baryon sector). On the other hand, the $1/N_c$ power of a diagram is given by:

$$\nu_{1/N_c} = \frac{1}{2}(n_\pi + n_B) + L - 1 + \sum_i n_i(\nu_{O_i} + 1 - b_i/2), \tag{8}$$

where n_π is the number of external pions of vertex i, and ν_{O_i} the $1/N_c$ order of the spin-flavor operator of that vertex. Since ν_{O_i} can be negative (due to factors of G^{ia} in vertices), one can think of individual diagrams with ν_{1/N_c} negative and violating large N_c consistency, requiring cancellation with other diagrams. Such a sum will have to respect the mentioned lower bound on the $1/N_c$ power corresponding to the sum of such diagrams. The explicit example of such cancellation in the axial currents at one-loop is given below. One interesting point is that the leading in $1/N_c$ order of terms in the Lagrangian are primarily determined by the order of the spin-flavor operators according to Eq.(6).

One can determine now the nominal counting of the one-loop contributions to the baryon masses and axial currents. The one loop correction shown in Fig. 1 has : $(L = 1, n_B = 2, n_\pi = 0, n_1 = 2, \nu_{O_1} = -1, b_1 = 2, \nu_{p_1} = 1)$ giving $\nu_p = 3$ as it is well known, and $\nu_{1/N_c} = -1$. Since there is only one possible diagram this must be consistent, and it is. For the axial currents one has the diagrams in Fig. IV. The current at tree level is $\mathcal{O}(N_c)$, and the sum of the diagrams cannot scale like a higher power of N_c. Performing the counting for the individual diagrams one obtains: $\nu_p(j) = 2$ for $j = 1, \cdots, 4$, and $\nu_{1/N_c}(j) = -2$, $j = 1, 2, 3$ and $\nu_{1/N_c}(4) = 0$. Thus a cancellation must occur of the $\mathcal{O}(N_c^2)$ terms when the contributions to the axial currents by diagrams 1, 2 and 3 are added. Since the acceptable bound is that the sum be $\mathcal{O}(N_c)$, one concludes that the axial current has, at one-loop, corrections $\mathcal{O}(p^2 N_c)$ or higher.

One can consider the case of two-loop diagrams. Consider diagrams where the same pion-baryon vertex Eq.(5) appears four times. For the masses one has $\nu_p(j) = 5$, and individual diagrams give $\nu_{1/N_c} = -2$. A cancellation must occur to restore the bound on the N_c counting for the masses, i.e., $\mathcal{O}(N_c)$. Thus, at two-loops the UV divergencies of the masses must be $\mathcal{O}(p^5 N_c)$ or higher. For the axial currents a similar discussion requires that counter-terms to the axial currents must be $\mathcal{O}(p^4 N_c)$ or higher. One loop with the term proportional to c_1 in Eq.(5) give $\mathcal{O}(\xi^4)$ contributions to the masses.

In order to determine the power counting in ξ for a given Feynman diagram with a single baryon flowing through the diagram, it is sufficient to know the $1/N_c$ power counting of the operators in the vertices, the number of external pions, and the chiral power counting of the vertices. Eliminating the numbers of pion and baryon propagators in the diagram using the known topological formulas [48]. A quick calculation then permits to express the order in ξ

of the diagram according to:

$$\nu_\xi = 1 + 3L + \frac{1}{2}n_\pi + \sum_i n_i(\nu_i + p_i - 1) \tag{9}$$

where L is the number of loops, n_π is the number of external pions, n_i the number of vertices of type i, and ν_i is the $1/N_c$ order of the spin-flavor operator appearing in and p_i the low energy (chiral) power of that type of vertex. It should be emphasized that because ν_i can be negative (due to N_c enhancement when the generator G^{ia} is present as a factor in a vertex). The ξ-power counting of the UV divergencies is obvious from the earlier discussion. At one-loop one finds that the masses have $\mathcal{O}(\xi^2)$ and $\mathcal{O}(\xi^3)$ counter-terms, while the axial currents will have $\mathcal{O}(\xi)$ and $\mathcal{O}(\xi^2)$ counter-terms. To two loops one expects $\mathcal{O}(\xi^4)$ and $\mathcal{O}(\xi^5)$, and $\mathcal{O}(\xi^3)$ and $\mathcal{O}(\xi^4)$ counter-terms for masses and axial currents respectively. The non-commutativity of limits is manifested in the finite terms where M_π and or momenta and δm appear combined in non-analytic terms, and are therefore sensitive to the linking of the two expansions.

III. BARYON MASSES

In this section baryon masses are analyzed to order ξ^3, or next-to-next to leading order (NNLO), in the limit of exact isospin symmetry. To that order the mass of the baryon of spin S reads:

$$m_{\mathbf{B}}(S) = N_c m_0 + \frac{C_{HF}}{N_c} S(S+1) + c_1 N_c M_\pi^2 + \delta m_{\mathbf{B}}^{1-loop+CT}(S), \tag{10}$$

where $\delta m_{\mathbf{B}}^{1-loop+CT}(S)$ involves contributions from the one-loop diagram in Fig. 1, and CT denotes counter-terms. From both types of contributions, there are $\mathcal{O}(\xi^2)$ and $\mathcal{O}(\xi^3)$ terms, and the calculation is exact at the latter order, as can be deduced from the previous discussion on power counting. The notation for the $\mathcal{O}(\xi)$ mass shift is used: $\delta m(S) \equiv \frac{C_{HF}}{N_c} S(S+1) + c_1 N_c M_\pi^2$. Notice that C_{HF} is equal to the LO term in $M_\Delta - M_N$ in the real world $N_c = 3$.

The leading 1-loop correction to the baryon self energy, diagram in Fig. 1, and reads:

$$\delta\Sigma_{(1-loop)} \;=\; i\,\frac{\mathring{g}_A^2}{F_\pi^2}\,\frac{1}{d-1}\,\sum_n G^{ia}\mathcal{P}_n G^{ia}\; I_{(1-loop)}(\delta m_n - p^0, M_\pi)\,, \tag{11}$$



FIG. 1: One-loop contribution to baryon self energy. The red propagator indicates sum over all possible baryons that can contribute.

where n indicates the possible intermediate baryon spin-isospin states in the loop, $\mathcal{P}_n$ are the corresponding spin-flavor projection operators, $\delta m_n = \delta m(S_n)$, and the loop integral is calculated in dimensional regularization with the result,

$$\begin{aligned} I_{(1-loop)}(Q, M_\pi) &= \int \frac{d^dk}{(2\pi)^d} \frac{\vec{k}^2}{k^2 - M_\pi^2 + i\epsilon} \frac{1}{k^0 - Q + i\epsilon} \\ &= \frac{i}{16\pi^2} \Bigg\{ Q \left((3M_\pi^2 - 2Q^2)(\lambda_\epsilon - \log \frac{M_\pi^2}{\mu^2}) + (5M_\pi^2 - 4Q^2) \right) \\ &+ 2\pi (M_\pi^2 - Q^2)^{3/2} + 4(Q^2 - M_\pi^2)^{3/2} \tanh^{-1} \frac{Q}{\sqrt{Q^2 - M_\pi^2}} \Bigg\} , \end{aligned} \tag{12}$$

where $Q = \delta m_n - p^0$, $\lambda_\epsilon = \frac{1}{\epsilon} - \gamma + \log 4\pi$, and μ is the renormalization scale which will be taken later to be of the order of m_ρ. For the specific evaluation of $\delta\Sigma_{(1-loop)}$ for a given baryon state denoted by in, $p^0 = \delta m_{in} - \mathfrak{p}^0$, where $\mathfrak{p}^0$ is a residual energy (when evaluated on an on-shell baryon it is the kinetic energy which is $\mathcal{O}(p^2/N_c)$). The non-commutativity of the $1/N_c$ expansion and the Chiral expansion of course resides in the non-analytic terms of the loop integral though their dependence on the ratio Q^2/M_π^2.

Appendix D provides all the necessary elements for the evaluation of the spin-flavor matrix elements in Eq. (11) as well as in the calculation of the one-loop corrections to the axial currents below.

The one-loop contribution to the wave function renormalization constant is given by:

$$\delta Z_{1-loop} = \frac{\partial}{\partial \mathfrak{p}^0} \delta\Sigma_{(1-loop)} \Big|_{\mathfrak{p}^0 \to 0} . \tag{13}$$

The explicit evaluation of the ultraviolet divergent pieces of the self energy gives:

$$\begin{aligned} \delta\Sigma^{UV}_{(1-loop)} &= \frac{\lambda_\epsilon}{16\pi^2} \frac{\mathring{g}_A^2}{F_\pi^2} \\ \times & \left\{ \frac{C_{HF}}{24 N_c} \left(-3M_\pi^2 (3N_c(4+N_c) - 20\hat{S}^2) + 8\frac{C_{HF}^2}{N_c^2}(N_c(4+N_c)(3+5\hat{S}^2) - 4\hat{S}^2(6+7\hat{S}^2)) \right) \right. \\ + & \left. \mathfrak{p}^0 \left(\frac{M_\pi^2}{2} (\frac{3}{8} N_c(4+N_c) - \hat{S}^2) - \frac{C_{HF}^2}{4N_c^2}(N_c(4+N_c)(3+2\hat{S}^2) - 8\hat{S}^2(3+\hat{S}^2)) \right) + \mathcal{O}(\mathfrak{p}^{0^2}) \right\} . \end{aligned} \tag{14}$$

The UV divergent pieces start at $\mathcal{O}(\xi^2)$. Note that the UV divergencies in the mass (term independent of $\mathfrak{p}^0$) is produced by the contribution of the partner baryon and is proportional to the mass splitting. As is well known they are absent in HBChPT without explicit Δ. The $\mathcal{O}(N_c^0)$ UV divergence is spin-flavor singlet and proportional to M_π^2, while the contributions to mass splittings are $\mathcal{O}(1/N_c^2)$. Notice that the leading UV divergence of δZ_{1-loop} is $\mathcal{O}(M_\pi^2 N_c)$: this is necessary as shown later for rendering the one-loop calculation of the axial currents consistent in the large N_c limit. Since the calculation is accurate to $\mathcal{O}(\xi^3)$, additional terms in the effective Lagrangian up to that order are necessary for renormalization. The terms necessary for renormalizing the self energy are therefore the following:

$$\begin{aligned}\mathcal{L}_\Sigma^{CT} &= \mathbf{B}^\dagger \left\{ \frac{m_1(N_c)}{N_c} + \frac{C_{HF1}(N_c)}{N_c^2}\hat{S}^2 + \frac{C_{HF2}(N_c)}{N_c^3}\hat{S}^4 + \mu_1(N_c)\chi_+ + \frac{\mu_2(N_c)}{N_c}\chi_+\hat{S}^2 \right. \\ &+ \left. \left(\frac{w_1(N_c)}{N_c} + \frac{w_2(N_c)}{N_c}\hat{S}^2 + \frac{w_3(N_c)}{N_c^3}\hat{S}^4 + (z_1(N_c)N_c + \frac{z_2(N_c)}{N_c}\hat{S}^2)\chi_+ \right)(iD_0 - \delta m) \right\} \mathbf{B}\end{aligned} \quad (15)$$

where one identifies the residual energy $\mathfrak{p}^0$ with the operator $(iD_0 - \delta m)$. All LECs are here of the form $X(N_c) = X_0 + X_1/N_c + \cdots$. Writing $X = X(\mu) + \gamma_X \lambda_\epsilon$, one renormalizes the self energy to $\mathcal{O}(\xi^3)$. The coefficients γ_X are determined from $\delta\Sigma^{UV}_{(1-loop)}$ given above. While the counter-terms are defined such that $X(\mu)$ is $\mathcal{O}(N_c^0)$, it is possible that γ_X is of higher order in $1/N_c$. Notice that among the higher order terms there are terms which can be simply absorbed into $1/N_c$ corrections to the LECs of the lowest order Lagrangian, and to m_0.

Finally, the baryon masses are given by:

$$m_{\mathbf{B}} = N_c m_0 + \frac{C_{HF}}{N_c}\hat{S}^2 + c_1 N_c M_\pi^2 + (\delta\Sigma^{UV\ finite}_{(1-loop)} + \delta\Sigma^{CT})|_{\mathfrak{p}^0=0}\,(1 + \delta Z^{UV\ finite}_{1-loop} + \delta Z^{CT}) \quad (16)$$

Note that the correction to the wave function renormalization factor enters in the expression for the mass corrections: this is because $\delta\Sigma(\mathfrak{p}^0 = 0)$ starts with terms $\mathcal{O}(\xi^2)$ and δZ starts at $\mathcal{O}(\xi)$, therefore the $\mathcal{O}(\xi^3)$ terms of the mass correction involve these lower order terms of the wave function renormalization.

The one-loop corrections and corresponding counter-terms contribute to the masses at $\mathcal{O}(\xi^2)$ and $\mathcal{O}(\xi^3)$, while in a strict large N_c limit the following ordering is found:

$$\begin{aligned} M_B &= \mathcal{O}(N_c) + \mathcal{O}(N_c M_\pi^2) + \mathcal{O}(N_c^0 M_\pi) + \cdots \\ M_B - M_B' &= \mathcal{O}(\frac{1}{N_c}) + \mathcal{O}(\frac{1}{M_\pi N_c^2}) \end{aligned} \quad (17)$$

Obviously the term $\mathcal{O}(\frac{1}{M_\pi N_c^2})$ stems from the $1/N_c$ expansion of non-analytic terms and shows the non-commutativity of limits.

The σ-terms for N and Δ, defined by $\sigma_B = \hat{m}\frac{\partial m_B}{\partial \hat{m}}$ ($\hat{m} = \frac{1}{2}(m_u + m_d)$), $\mathcal{O}(\xi)$ with the one-loop corrections contributing up to $\mathcal{O}(\xi^3)$. The difference $\sigma^B - \sigma^{B'}$ is $\mathcal{O}(\xi^2)$ and at that order it receives only finite contributions from the loop. This implies that the slopes of the N and Δ masses as functions of M_π are the same up to $\mathcal{O}(\xi^2)$ deviations. This seems to be closely followed by the lattice QCD results analyzed later. In the large N_c limit, obviously $\sigma = \mathcal{O}(N_c)$. The terms of that order are necessarily spin-flavor singlet, and taking the limit at fixed M_π one finds $\sigma_\Delta - \sigma_N = \mathcal{O}(1/N_c^2)$, a result similar to the one in the ξ-expansion.

IV. AXIAL COUPLINGS

In this section the evaluation of the axial couplings including corrections $\mathcal{O}(\xi^2)$ is presented. At that order the one-loop corrections must be calculated.

The matrix elements of interest for the axial currents are $\langle \mathbf{B'} \mid A^{ia} \mid \mathbf{B}\rangle$ evaluated at vanishing external 3-momentum. The axial couplings are then defined by:

$$\langle \mathbf{B'} \mid A^{ia} \mid \mathbf{B}\rangle = g_A^{\mathbf{BB'}}\,\frac{5}{6}\,\langle \mathbf{B'} \mid G^{ia} \mid \mathbf{B}\rangle\,. \tag{18}$$

The axial couplings defined here are $\mathcal{O}(N_c^0)$. The $\mathcal{O}(N_c)$ of the matrix elements of the axial currents is due to the operator G^{ia}. The factor 5/6 mentioned earlier is included so that g_A^{NN} at $N_c = 3$ exactly corresponds to the usual nucleon g_A, which has the value 1.2701 ± 0.0025 [49]. This definition of the axial couplings is convenient in the context of the $1/N_c$ expansion, as the differences between the different axial couplings are $\mathcal{O}(1/N_c^2)$.



FIG. 2: Diagrams contributing to the 1-loop corrections to the axial-currents. The crossed circle denotes the axial-current operator.

The determination of the axial couplings to $\mathcal{O}(\xi^2)$ require the calculation of the 1-loop corrections to the axial current. Only the contributions with no pion pole are necessary, and they are given by the diagrams in Fig. IV. The resulting 1-loop contribution to the axial currents reads:

$$\delta A^{ia}_{1-loop} = \delta A^{ia}_{1-loop}(1) + \delta A^{ia}_{1-loop}(2+3) + \delta A^{ia}_{1-loop}(4)\,., \tag{19}$$

where $\delta A^{ia}_{1-loop}(2+3)$ is given by a factor 1/2 times the no-baryon-pole contributions of diagrams (2+3). The different contributions read as follows, where one needs to take the limits $\mathfrak{p}^0, \mathfrak{p}'^0 \to 0$:

$$\begin{aligned}
\delta A^{ia}_{1-loop}(1) &= -i\frac{\mathring{g}_A^3}{F_\pi^2}\frac{1}{d-1} \\
&\quad\times \sum_{n,n'} G^{jb}\mathcal{P}_{n'}G^{ia}\mathcal{P}_n G^{jb}\frac{I_{1-loop}(\delta m_n - p^0, M_\pi) - I_{1-loop}(\delta m_{n'} - p'^0, M_\pi)}{p^0 - p'^0 - \delta m_n + \delta m_{n'}} \\
\delta A^{ia}_{1-loop}(2+3) &= \frac{\mathring{g}_A}{2}\left(G^{ia}\delta Z_{1-loop} + \delta Z_{1-loop}G^{ia}\right) \\
\delta A^{ia}_{1-loop}(4) &= -\frac{\mathring{g}_A}{3F_\pi^2}\Delta(M_\pi)G^{ia}.
\end{aligned} \tag{20}$$

Obviously, G^{ia} and δZ_{1-loop} do not commute in general. The pion tadpole integral in the last term is given by:

$$\Delta(M_\pi) = -\frac{M_\pi^2}{16\pi^2}(\lambda_\epsilon - \log\frac{M_\pi^2}{\mu^2}). \tag{21}$$

Notice that the contribution by diagram (4) is actually $\mathcal{O}(\xi^4)$, and thus beyond the degree of accuracy of the present calculation. It can serve however as a measure of the size of the NNNLO corrections.

The corrections to the axial currents must scale as $\mathcal{O}(N_c^\nu)$ with $\nu \leq 1$. While diagram (4) is $\mathcal{O}(N_c^0)$ and therefore consistent in itself, diagrams (1) and (2+3) above are $\mathcal{O}(N_c^2)$. As shown in Ref. [28], the offending terms cancel upon adding the diagrams. To test the cancellation it is sufficient to take the large N_c limit at fixed M_π. A straightforward evaluation leads to:

$$\delta A^{ia}_{1-loop}(1)+\delta A^{ia}_{1-loop}(2+3)\Big|_{N_c\to\infty} = -i\frac{\mathring{g}_A^3}{F_\pi^2}\frac{1}{d-1}\left\{\frac{1}{2}[[G^{jb},G^{ia}],G^{jb}]\frac{\partial}{\partial\mathfrak{p}^0}I_{1-loop}(\mathfrak{p}^0,M_\pi)+\cdots\right\}, \tag{22}$$

where $\cdots$ indicate further terms which are consistent with the N_c power counting of the axial currents. The suppression of the $\mathcal{O}(N_c^2)$ terms is direct consequence of the appearance of the commutator of two generators G, which is $\mathcal{O}(N_c^0)$, when the diagrams are added up. In consequence the displayed terms are $\mathcal{O}(N_c^0)$.

The UV divergent contributions of the individual diagrams read (using the notation

$\hat{S}^n \equiv (S(S+1))^n)$:

$$\begin{aligned}
\delta A^{ia}_{1-loop}(1)^{UV} &= -\frac{\lambda_\epsilon}{48\pi^2}\frac{\mathring{g}_A^3}{F_\pi^2}\Big\{3M_\pi^2 G^{jb}G^{ia}G^{jb} \\
&- 2\left(\frac{C_{HF}}{N_c}\right)^2 \Big(G^{jb}G^{ia}\left[\left[G^{jb},\hat{S}^2\right],\hat{S}^2\right] + \left[\hat{S}^2,\left[\hat{S}^2,G^{jb}\right]\right]G^{ia}G^{jb} \\
&+ \left[\hat{S}^2,G^{jb}\right]G^{ia}\left[G^{jb},\hat{S}^2\right]\Big)\Big\} \\
\delta A^{ia}_{1-loop}(2+3)^{UV} &= \frac{\lambda_\epsilon}{96\pi^2}\frac{\mathring{g}_A^3}{F_\pi^2}\Big\{3M_\pi^2\{G^{ia},G^2\} \\
&- 2\left(\frac{C_{HF}}{N_c}\right)^2\Big(G^{ia}G^{jb}[[G^{jb},\hat{S}^2],\hat{S}^2] + [\hat{S}^2,[\hat{S}^2,G^{jb}]G^{jb}G^{ia}\Big)\Big\} \\
\delta A^{ia}_{1-loop}(4)^{UV} &= \frac{\lambda_\epsilon}{48\pi^2}\frac{\mathring{g}_A}{F_\pi^2}\,M_\pi^2\,G^{ia}. \qquad (23)
\end{aligned}$$

One notices that only the terms proportional to M_π^2 in diagrams (1) and (2+3) diverge as proportional to N_c^2, while the terms proportional to C_{HF}^2 are $\mathcal{O}(N_c^0)$. Thus only the $\mathcal{O}(N_c^2)$ of the terms proportional to M_π^2 need to cancel to give consistency. One can easily check that such a cancellation indeed occurs, leaving only terms $\mathcal{O}(N_c^0)$. An explicit evaluation of these UV divergent terms using the results from Appendix D finally gives:

$$\begin{aligned}
\delta A^{ia}_{1-loop}{}^{UV} &= \frac{\lambda_\epsilon}{32\pi^2}\frac{\mathring{g}_A}{F_\pi^2 N_c^2}\Big\{(\frac{2}{3}+\mathring{g}_A^2)M_\pi^2 N_c^2 G^{ia} \\
&+ \frac{C_{HF}^2\mathring{g}_A^2}{3}\Big(4 - 2N_c(4+N_c)G^{ia} - 7[\hat{S}^2,[\hat{S}^2,G^{ia}]] + 4\{\hat{S}^2,G^{ia}\}\Big)\Big\} \qquad (24)
\end{aligned}$$

The terms in the Lagrangian needed to renormalize the axial currents are then the following:

$$\mathcal{L}_A^{CT} = \mathbf{B}^\dagger u_{ia}\left(\frac{C_0^A}{N_c}G^{ia} + \frac{C_1^A}{4}\{\chi_+,G^{ia}\} + \frac{C_2^A}{N_c^2}\{\hat{S}^2,G^{ia}\} + \frac{C_3^A}{N_c}[\hat{S}^2,G^{ia}] + \frac{C_4^A}{N_c}S^i I^a\right)\mathbf{B} \qquad (25)$$

These are all the terms which can contribute to the axial currents up to $\mathcal{O}(\xi)$, which will determine the axial couplings up to $\mathcal{O}(\xi^2)$, i.e., NNLO, which is what is needed for our purpose. There are several very important observations concerning the ξ-power counting. The corrections to the axial couplings start at $\mathcal{O}(\xi)$, and the individual contributions of the different baryons in the loop diagrams are also $\mathcal{O}(\xi)$. Even the difference of different axial couplings $g_A^{BB'} - g_A^{B''B'''}$ starts at $\mathcal{O}(\xi)$. These latter differences are UV finite. The large N_c cancellations do not seem manifest. However, at $N_c = 3$, where the ξ-expansion is used, cancellations do occur numerically as shown by Fig. 4 in Section V. Thus, the smallness of $\mathcal{O}(\xi)$ terms in the axial couplings is the incipient manifestation of the cancellations in the large N_c limit. If one would consider the strict large N_c limit, the one loop corrections and counter-terms considered give the following $1/N_c$ power counting:

$$\begin{aligned} g_A^{BB'} - \frac{5}{6}\mathring{g}_A &= \mathcal{O}(M_\pi^2) + \log\left(\frac{M_\pi^2}{\mu^2}\right)\mathcal{O}(\frac{M_\pi^2}{N_c}) + \cdots \\ g_A^{BB'} - g_A^{B''B'''} &= \mathcal{O}(\frac{1}{N_c^2}) + \mathcal{O}(\frac{M_\pi}{N_c^2}), \end{aligned} \tag{26}$$

where, as expected, the latter differences are UV finite as in the ξ expansion.

While in next section a discussion of the nucleon's g_A in the context of LQCD results, one can make an estimate of the spin-flavor symmetry breaking terms in the axial couplings g_A^{NN} vs $g_A^{\Delta N}$ using the result for the Δ width:

$$\Gamma_{\Delta\to\pi N} = \frac{1}{12\pi}\left(\frac{6}{5}\frac{g_A^{\Delta N}}{F_\pi}\right)^2((m_\Delta - m_N)^2 - M_\pi^2)^{3/2} \tag{27}$$

with the experimental value $\Gamma_{\Delta\to\pi N}(Exp) = 116 - 120$ MeV [49], one obtains $g_A^{\Delta N} = 1.235 \pm 0.011$, which is remarkably close to $g_A^{NN} = 1.2701 \pm 0.0025$ [49].

V. ANALYSIS OF LATTICE QCD RESULTS FOR BARYON MASSES AND THE NUCLEON'S AXIAL COUPLING

As an application of the present framework of the ξ-expansion, this section presents an analysis of LQCD results for baryon masses and the nucleon's axial coupling.

Lattice QCD calculations of the non-strange ground state baryon masses (both of N and Δ baryons) have opened the possibility of determining the quark mass dependencies, and similarly for the axial coupling of the nucleon. These calculations represent a very fruitful ground of applications for ChPT, allowing in particular for a study of the convergence of the low energy expansion. Modern dynamical three-flavor ($N_f = 2 + 1$) calculations of the hadron spectrum, and in particular of baryon masses, with fixed strange quark mass and variable $m_u = m_d$ [34–39] are achieving remarkably accurate results in a range of quark masses where extrapolations to the physical limit are now possible using effective theory. All calculations present similar results for the N and Δ masses, namely, roughly linear dependencies of the masses as a function of M_π, and extrapolations to the correct physical value within a few percent. For the nucleon axial coupling g_A^{NN} the results are particularly interesting [40–45] because it shows small dependence in a broad range of M_π. The most recent LQCD calculations for $N_f = 2$ [40, 42, 45] and $N_f = 2+1$ [41, 43, 44], all agree on that observation. An open issue is that all calculations give an underestimation for the value of g_A^{NN} of about 12% below the experimental value.

Effects due to finite volume of the lattice have been studied for the observables considered here. Those effects are determined primarily by the value of the product LM_π, where L is the length of the lattice. For the baryon masses, the rule $LM_\pi \sim 4$ [33] seems to be sufficient for the volume effects to be negligibly small. On the other hand, for g_A the LQCD understanding of the finite volume effects is not yet complete. According to Ref. [43], g_A clearly exhibits scaling in LM_π and in lattices with $LM_\pi \sim 4-5$ the effect on g_A is a 9 % reduction in calculations with $2+1$ flavors of domain wall fermions and a 25 % reduction in calculations with 2 flavors of Wilson fermions. This has led to the current view that $LM_\pi \gtrsim 5-6$ or even higher may in fact be needed to reliably determine g_A. Finite-volume effects for mases and the nucleon axial coupling have been studied in effective theories [50–59]. A detailed study of these effects in the present formalism is beyond the scope of this work, and will be presented elsewhere [60].

In the following, combined fits to LQCD results for N and Δ masses and the nucleon g_A as functions of M_π are carried out. Several studies of the available results have been carried out, but only the analysis of the more recent ones are discussed here. For the N and Δ masses the results used are those from the PACS-CS collaboration of Ref. [36] and the LHP collaboration of Ref. [35]. For g_A the results used are those from the LHP collaboration [41] and from the ETM collaboration [42]. All collaborations obtain results satisfying the constraint $LM_\pi \gtrsim 4$ and for quark masses reaching down close to the physical point, in particular for the baryon masses. The fits are carried out only including results where $LM_\pi \gtrsim 4$. The analysis of these LQCD results is carried out up to $\mathcal{O}(\xi^3)$ for the masses and $\mathcal{O}(\xi^2)$ for g_A. The set of Lagrangian counter-terms is the one displayed in Eqs. (16) and (25), which are summarized by the following equation:

$$\begin{aligned}
\delta\Sigma^{CT}(\mathfrak{p}^0=0)(S) &= \frac{m_1}{N_c} + \frac{C_{HF1}}{N_c^2}\hat{S}^2 + \frac{C_{HF2}}{N_c^3}\hat{S}^4 + \mu_1 M_\pi^2 + \frac{\mu_2}{N_c}\hat{S}^2 M_\pi^2\,,\\
\delta Z^{CT}(S) &= \frac{w_1}{N_c} + \frac{w_2}{N_c}\hat{S}^2 + \frac{w_3}{N_c^3}\hat{S}^4 + z_1 N_c M_\pi^2 + \frac{z_2}{N_c}\hat{S}^2 M_\pi^2\,,\\
\delta g_A^{CT}(S,S') &= \frac{C_0^A}{N_c} + C_1^A M_\pi^2 + \frac{C_2^A}{N_c^2}(\hat{S}^2+\hat{S}'^2) + \frac{C_3^A}{N_c^2}(\hat{S}^2-\hat{S}'^2)\\
&+ \frac{4C_4^A}{N_c(2+N_c)}\delta_{SS'}\hat{S}^2\,.
\end{aligned} \tag{28}$$

There are several LECs which to be determined require knowledge of results at different values of N_c. With the LQCD results at fixed $N_c = 3$, those LECs combine with existing ones at lower order, making their determination impossible. In addition, because LQCD results on $g_A^{N\Delta}$ are not analyzed and the lack of results for $g_A^{\Delta\Delta}$, LECs which split the values of the different g_A's cannot be fixed either. For instance, the LECs m_1 and w_1 give the sub-leading N_c dependence m_0, and therefore at fixed $N_c = 3$ they are absorbed into the fitted value of m_0. The same will happen with C_{HF1} and w_2 with C_{HF}. The LEC μ_1 is a correction to the LO σ-term LEC c_1. Similarly, one can not separate C_0^A from $\mathring{g}_A$. Therefore, without loss of generality, the redundant LECs can be set to vanish. In addition, since the current fits only involve the nucleon's axial coupling, g_A^{NN}, not all LECs affecting the axial currents can be determined as mentioned earlier. In particular, counter-terms with commutators in Eq. (25) only appear in $g_A^{N\Delta}$. Of course, depending on the order in the ξ-expansion, the number of LECs varies. Specifically, at leading order (LO), that is $\mathcal{O}(\xi)$ for the mass and $\mathcal{O}(\xi^0)$ for the axial coupling, the LECs are m_0, $\mathring{g}_A$, C_{HF} and c_1, at NLO the additional LECs C_{HF1}, μ_1 and C_0^A appear. Finally, at NNLO, that is $\mathcal{O}(\xi^3)$ for the mass and $\mathcal{O}(\xi^2)$ for the axial coupling, the additional LECs μ_2, z_1 and C_1^A, which are fitted, and μ_3, w_1, w_2 and $C_{2,3,4}^A$ that cannot be determined, make their appearance.

The combined fits to N and Δ masses and to g_A^{NN} up to NNLO for the four possible combinations of LQCD results from the collaborations considered here are presented in table I, which shows the values for LECs obtained from the fits and the extrapolated values for m_N, m_Δ and g_A to the physical point. To estimate the theoretical errors, the original lattice results are bootstrapped by Montecarlo, and the errors correspond to a 68% confidence interval. The LO fits are restricted to pion masses in the range $M_\pi \sim 140 - 300$ MeV, NLO fits to $M_\pi \sim 140 - 500$ MeV, and NNLO fits to $M_\pi \sim 140 - 600$ MeV.

The following remarks on the fits are in order:

1. All fitted LECs are of natural size when the renormalization scale is taken to be $\mu \sim m_\rho$.

2. Parameters appearing at lower orders, namely m_0, $\mathring{g}_A$, C_{HF} remain stable at higher orders, except c_1 which changes by more than the estimated 30% when increasing the order in ξ of the fit by one unit.

3. For baryon masses, LQCD data and physical point values are consistent even at LO

Collab.	Order	m_0 [MeV]	C_{HF} [MeV]	$\mathring{g}_A$	c_1 [MeV^{-1}]	μ_2 [MeV^{-1}]	z_1 (10^{-7}) [MeV^{-2}]	C_1^A (10^{-7})	m_N [MeV]	m_Δ [MeV]	g_A^{NN}	χ^2	χ^2/DOF
PACS-CS + LHP	LO	271(8)	301(19)	1.381(11)	0.0008(1)	N/A	N/A	N/A	935(17)	1236(17)	1.15(1)	20.15	3.36
	NLO	274(7)	208(9)	1.391(12)	0.00133(5)	N/A	N/A	N/A	957(14)	1222(14)	1.13(1)	28.80	3.20
	N^2LO	264(7)	169(9)	1.427(16)	0.0023(1)	-0.0002(1)	-9.0(5)	-1.7(6)	957(14)	1195(14)	1.131(12)	41.79	3.48
LHP + LHP	LO	264(6)	378(13)	1.381(11)	0.00075(5)	N/A	N/A	N/A	928(14)	1306(15)	1.15(1)	59.69	9.95
	NLO	267(4)	230(5)	1.391(11)	0.00143(2)	N/A	N/A	N/A	961(9)	1257(10)	1.13(1)	73.08	8.12
	N^2LO	223(5)	213(6)	1.366(14)	0.00255(6)	-0.00046(6)	-2.7(2)	-1.8(0.5)	886(10)	1180(12)	1.08(1)	147.97	12.33
PACS-CS + ETM	LO	272(8)	301 (20)	1.395(15)	0.0008(1)	N/A	N/A	N/A	936(17)	1236(17)	1.162(13)	21.01	2.10
	NLO	274(7)	208(8)	1.399(13)	0.00134(5)	N/A	N/A	N/A	958(15)	1222(14)	1.139(11)	29.93	2.30
	N^2LO	263(7)	166(9)	1.47(3)	0.0024(1)	-0.00023(11)	-9.3(6)	-3.7(1.6)	957(14)	1192(14)	1.158(19)	42.89	2.68
LHP + ETM	LO	264(6)	379(12)	1.395(15)	0.00075(5)	N/A	N/A	N/A	928(14)	1307(14)	1.162(12)	60.55	6.06
	NLO	267(4)	230(5)	1.399(13)	0.00144(3)	N/A	N/A	N/A	960(9)	1256(10)	1.137(11)	74.06	5.69
	N^2LO	227(5)	217(7)	1.326(18)	0.00241(7)	-0.00047(6)	-2.7(2)	3.5(1.2)	895(10)	1193(12)	1.055(12)	143.13	8.95
Natural magnitude of LECs		~ 300	~ 300	1-2	10^{-3}	10^{-3}	10^{-6}	10^{-6}					

TABLE I: Results for LECs and extrapolated masses m_N and $Re(m_\Delta)$, and the nucleon g_A from a combined fit to LQCD. The renormalization scale has been taken to be $\mu = 700$ MeV. Lattice results for masses are from the PACS-CS [36] and LHP [35] collaborations, and for g_A from LHP [35, 41] and ETM [42] collaborations. Different orders in the ξ-expansion are shown. The LECs not shown have been set to vanish.



FIG. 3: Combined fits to PACS-CS [36] and LHP [41] corresponding to the results shown in the first row of Table I. LO (black long-dashed line), NLO (black short-dashed line) and NNLO for N (blue solid line) and Δ (red solid line). The bands correspond to the theoretical 68% confidence interval.

where only with three parameters one can extrapolate to the correct experimental values. This is not surprising taking into account that a single straight line can be use for consistent fit [35].

4. For the case of the axial current, cancellations of large contributions from individual loop diagrams are very pronounced and the almost flat behavior of g_A as a function of M_π obtained in LQCD is naturally explained. This is shown in the left panel of Fig. 4 which shows the finite one-loop contributions to g_A^{NN} from each diagram ($\mu = 700$ MeV). As stated in Eq. (22) this cancellation is exact in the large N_c limit. However, at $N_c = 3$ this cancellation is not exact but still quite pronounced (black curve in Fig. 4), and plays the key role in explaining the small dependence in M_π. A similar cancellation occurs between the contributions of N and Δ in the loop. This is shown in the right panel of Fig. 4.

5. The physical g_A^{NN} cannot be fitted along with the lattice results, instead the lattice results and the expansion to NNLO extrapolate to a value 12% smaller value than the physical one. The recent LQCD results [61] which reach further down in M_π continue that trend. This seems to be therefore an issue with the LQCD calculations rather than the effective theory. In fact, it would be very unnatural for the effective theory



FIG. 4: Finite parts of the one-loop contributions to g_A^{NN}. The upper left panel shows the individual contributions of the diagrams in Fig. 2 up to $\mathcal{O}(\xi^3)$. The right panel shows the corresponding effect of switching off the contribution of the Δ in the loops. The third panel shows the effect of removing the contributions of the counter-terms to the masses. Throughout $\mu = 700$ MeV.

to describe a relatively fast changing pattern at low M_π, to continue with the slow changing behavior at larger M_π, which it can describe in a natural fashion.

6. A fit restricted only to masses gives too small a value for g_A^{NN}, namely, $g_A \sim 0.5 - 0.8$. A realistic value can only be obtained with the combined fit.

7. Predictions for $g_A^{N\Delta}$ and $g_A^{\Delta\Delta}$ cannot be made without the corresponding LQCD results. However, the results at the physical point from Eq. (27) suggest that these are going to be very similar in value to g_A^{NN}. Further efforts to study this couplings in LQCD will be very useful.

8. In the masses one finds that above $M_\pi > 350 - 400$ MeV there is a significant cancellation between the contributions of the one-loop diagram and the counter-terms as shown in Fig 4, which must be taken as an indicator of the range of convergence of the expansion. The masses counter-terms are $\mathcal{O}(M_\pi^2 N_c^0) = \mathcal{O}(\xi^2)$.

9. Using the results of the fit at $\mathcal{O}(\xi^3)$, one obtains for the σ terms evaluated at the physical pion mass:

$$\begin{aligned} \sigma^N &= 65 \pm 8\text{MeV} \\ \sigma^\Delta &= 90 \pm 9\text{MeV}. \end{aligned} \tag{29}$$

10. It must be emphasized that the results obtained here are very similar to those obtained in works where the Δ has been included explicitly [16, 19, 20, 62–64]. The main advantage of the present approach of the ξ-expansion is its systematic character, which in particular will be more prominently shown when carrying out higher order calculations than the ones considered here.

VI. DISCUSSION AND CONCLUSIONS

Chiral symmetry and the large N_c limit are of fundamental conceptual importance in QCD. The former is known to play a crucial role in light hadrons, and there are multiple indications that the latter is also important, in particular for baryons. It is therefore very important to have a theoretical framework where both of these aspects of QCD are consistently incorporated. This is possible with the combined $1/N_c$ and Chiral expansions of QCD, which in the baryon sector is implemented with the effective theory discussed in this work. A particular power counting, the ξ-expansion, which links the $1/N_c$ and low energy expansions as $1/N_c = \mathcal{O}(\xi) = \mathcal{O}(p)$ is proposed as the most realistic one for studying baryons at $N_c = 3$. Results for the masses and axial couplings at NNLO have been given, and applied to current LQCD results.

The ξ-expansion at NNLO clearly provides a satisfactory description of the LQCD results, and in particular it illuminates the mild dependence of the axial couplings on the quark masses as a result of important cancellations, which had been realized in various previous analysis by various groups. It is important to complete the study in $SU(3)$, in particular because the one-loop contributions to the baryon masses become larger in magnitude, and a smaller range of convergence is expected. These results will be presented elsewhere [65]

The deficit in g_A at the physical point seems to be a LQCD issue rather than a problem of convergence of the effective theory. In particular because the ξ-expansion is especially well behaved for g_A. It is however a puzzling problem, because the likely candidate for the explanation, namely finite volume effects, is expected to be significantly somewhat than the effect observed.

In addition to the tests LQCD can provide on the quark mass dependencies, it is the ideal tool to test the N_c behavior of QCD. Baryon LQCD is becoming accessible at varying values of N_c [66], which is a promising development.

Acknowledgments

The authors thank C. Alexandrou, K. Kanaya and D. Renner for useful discussions on several aspects of the LQCD results. This work was supported by DOE Contract No. DE-AC05-06OR23177 under which JSA operates the Thomas Jefferson National Accelerator Facility, and by the National Science Foundation (USA) through grant PHY-0855789 (JLG).

Appendix A: Spin-flavor Algebra

The $4N_f^2-1$ generators of the spin-flavor group $SU(2N_f)$ consist of the three spin generators S^i, the N_f^2-1 flavor $SU(N_f)$ generators T^a, and the remaining $3(N_f^2-1)$ spin/flavor generators G^{ia}. The commutation relations are:

$$\begin{gathered}[S^i,S^j]=i\epsilon_{ijk}S^k,\quad [T^a,T^b]=if_{abc}T^c,\quad [T^a,S^i]=0\\ [S^i,G^{ja}]=i\epsilon_{ijk}G^{ka},\quad [T^a,G^{ib}]=if_{abc}G^{ic}\\ [G^{ia},G^{jb}]=\tfrac{i}{4}\delta^{ij}f^{abc}T^c+\tfrac{i}{2N_f}\delta^{ab}\epsilon^{ijk}S^k+\tfrac{i}{2}\epsilon^{ijk}d^{abc}G^{kc}\end{gathered} \tag{A1}$$

For two flavors one has the isospin generators I^a $a=1,2,3$.

In representations with N_c indices (baryons), the generators G^{ia} have matrix elements $\mathcal{O}(N_c)$ on states with $S=\mathcal{O}(N_c^0)$. A contracted $SU(4)$ algebra is defined by the generators $\{S^i,I^a,X^{ia}\}$, where $X^{ia}=G^{ia}/N_c$. In large N_c, the generators X^{ia} become semiclassical as $[X^{ia},X^{jb}]=\mathcal{O}(1/N_c^2)$, while having matrix elements $\mathcal{O}(1)$ in baryon representations.

Appendix B: Non-linear realization of chiral symmetry and spin-flavor transformations

In the symmetric representations of $SU(4)$ the baryon spin-flavor multiplet consists of the baryon states with $I=S$. In particular, isospin transformation will act on the spin-flavor multiplet in an obvious way on those states. This permits a straighforward implementation of the non-linear realization of chiral $SU_L(2)\times SU_R(2)$ on the spin-flavor multiplet. Defining as usual the Goldstone Boson fields π^a through the unitary parametrization $u=\exp(i\frac{\pi^a I^a}{2F_\pi})$, for any isospin representation one defines a non-linear realization of chiral symmetry according to [3, 4]:

$$(L,R):u=u'=Ruh^\dagger(L,R,u)=h(L,R,u)uL^\dagger, \tag{B1}$$

where (L,R) is any $SU_L(2)\times SU_R(2)$ transformation. This equation defines h, and since h is an isospin $SU(2)$ transformation itself, it can be written as $h=\exp(ic^aI^a)$. The chiral transformation on the baryon multiplet $\mathbf{B}$ is then given by:

$$(L,R):\mathbf{B}=\mathbf{B}'=h(L,R,u)\mathbf{B}. \tag{B2}$$

On the other hand, spin-flavor transformations of interest are the contracted ones, namely those generated by $\{S^i,I^a,X^{ia}=\frac{1}{N_c}G^{ia}\}$. While the isospin transformations act on the pion fields in the usual way, and the spin transformations must be performed along with the corresponding spatial rotations, and the transformations generated by X^{ia} are defined to only act on the baryons.

Appendix C: Tools for building effective Lagrangians

The effective baryon Lagrangian can be expressed in the usual way as a series of terms which are $SU_L(2) \times SU_R(2)$ invariant (upon introduction of appropriate sources; see for instance [67] for details). In addition, implemented in the effective Lagrangian is the approximate $SU(4)$ symmetry and its breaking as a power series in $1/N_c$ [27]. The fields in the effective Lagrangian are the Goldstone Bosons parametrized by the unitary $SU(2)$ matrix field u and the baryons given by the symmetric $SU(4)$ multiplet $\mathbf{B}$ of $I = S$ fields.

The building blocks for the effective theory consist of low energy operators, and spin-flavor operators.

The low energy operators are the usual ones, namely:

$$\begin{aligned}
D_\mu &= \partial_\mu - i\Gamma_\mu, \quad \Gamma_\mu = \Gamma_\mu^\dagger = \frac{1}{2}(u^\dagger(i\partial_\mu + r_\mu)u + u(i\partial_\mu + \ell_\mu)u^\dagger) \\
u_\mu &= u_\mu^\dagger = u^\dagger(i\partial_\mu + r_\mu)u - u(i\partial_\mu + \ell_\mu)u^\dagger \\
\chi &= 2B_0(s + ip), \quad \chi_\pm = u^\dagger \chi u^\dagger \pm u\chi^\dagger u \\
F_L^{\mu\nu} &= \partial^\mu \ell^\nu - \partial^\nu \ell^\mu - i[\ell^\mu, \ell^\nu], \quad F_R^{\mu\nu} = \partial^\mu r^\nu - \partial^\nu r^\mu - i[r^\mu, r^\nu] \qquad \text{(C1)}
\end{aligned}$$

where D_μ is the chiral covariant derivative, s and p are scalar and pseudo-scalar sources, $\chi_\pm = 2M_\pi^2 + \cdots$, and ℓ_μ and r_μ are gauge sources. The spin-flavor operators are tensor operators consisting of products of the spin-flavor generators. These operators can be reduced by means of the commutation relations to forms which only contain anti-commutators. A set of identities shown in Table nnnn permits one to arrive at sets of basis operators at each order in $1/N_c$ for a given spin/isospin tensor type of operator. The $1/N_c$ order ν of an operator reduced as mentioned is $\nu = n - 1 - \kappa$, where n is the number of generators appearing as factors in the operator (one then says that the operator is an n-body operator), and κ is the number of generators G^{ia} in the product.

The leading order equations of motion can be used in the construction of the higher order terms, namely, $iD_0\mathbf{B} = (\frac{C_{HF}}{N_c}S(S+1) + \frac{c_1}{2}N_c\chi_+)\mathbf{B}$, and $\nabla_\mu u^\mu = \frac{i}{2}\chi_-$.

Appendix D: Matrix elements of spin-flavor operators in the symmetric representations of SU(4)

The evaluation of the matrix elements of spin-flavor operators in the present work can be carried out starting from the following matrix elements of the spin-flavor generators in the totally symmetric representation of $SU(4)$ corresponding to the Young tableux with a single row of N_c boxes. The basis states of the symmetric representation consists of the states with $I = S$, namely $\mid S\, S_3 I_3\rangle$, where S_3 and I_3 are the spin and isospin projections respectively.

$$\begin{aligned}
\langle S'\, S_3' I_3' \mid S^i \mid S\, S_3 I_3\rangle &= \sqrt{S(S+1)}\delta_{SS'}\delta_{I_3 I_3'}\langle S\, S_3, 1i \mid S'\, S_3'\rangle \\
\langle S'\, S_3' I_3' \mid I^a \mid S\, S_3 I_3\rangle &= \sqrt{S(S+1)}\delta_{SS'}\delta_{S_3 S_3'}\langle S\, I_3, 1a \mid S' I_3'\rangle \\
\langle S'\, S_3' I_3' \mid G^{ia} \mid S\, S_3 I_3\rangle &= \frac{1}{4}\sqrt{\frac{2S+1}{2S'+1}}\zeta(N_c, S, S')\langle S\, S_3, 1i \mid S'\, S_3'\rangle\langle S\, I_3, 1a \mid S' I_3'\rangle, \qquad \text{(D1)}
\end{aligned}$$

where $\zeta(N_c, S, S') = \sqrt{(2+N_c)^2 - (S-S')^2(S+S'+1)^2}$ [68]. The products of generators can be reduced by means of the use of the commutation relations, and further, for matrix elements in the symmetric representation, via the reduction rules [69], which for convenience are displayed in the following Table II.

TABLE II: $SU(4)$ operator identities in the totally symmetric irreducible representation $(N_c, 0, 0)$ of $SU(4)$. The last column gives the operator's quantum numbers (J, I) under $SU(2) \times SU(2)$

$\{S^i, S^i\} - \{I^a, I^a\} = 0$	(0,0)
$\{S^i, S^i\} + \{I^a, I^a\} + 4\{G^{ia}, G^{ia}\} = \frac{3}{2}N_c(4+N_c)$	(0,0)
$2\{S^i, G^{ia}\} = (2+N_c)I^a$	(0,1)
$2\{I^a, G^{ia}\} = (2+N_c)S^i$	(1,0)
$\frac{1}{2}\{S^k, I^c\} - \epsilon^{ijk}\epsilon^{abc}\{G^{ia}, G^{jb}\} = (2+N_c)G^{kc}$	(1,1)
$\epsilon^{ijk}\{S^i, G^{jc}\} = \epsilon^{abc}\{I^a, G^{kb}\}$	(1,1)
$4\{G^{ia}, G^{ib}\}\vert_{I=2} = \{I^a, I^b\}\vert_{I=2}$	(0,2)
$4\{G^{ia}, G^{ja}\}\vert_{J=2} = \{S^i, S^j\}\vert_{J=2}$	(2,0)

1. Useful matrix elements

It is always convenient to express matrix elements in terms of reduced matrix elements (RMEs) defined in the ordinary Wigner-Eckart fashion [70]. The RMEs defined here are with respect to $SU_{\rm spin}(2)\times SU_I(2)$. For matrix elements in the symmetric representation of spin flavor the Wigner-Eckart theorem reads:

$$\langle S'\, S'_3 I'_3 \mid \mathbf{O}^{\mathbf{JJ_3}}_{\mathbf{II_3}} \mid S\, S_3 I_3\rangle = \frac{\langle S' \mid\mid \mathbf{O}^{\mathbf{J}}_{\mathbf{I}} \mid\mid S\rangle}{2S'+1}\langle S S_3, \mathbf{JJ_3} \mid S' S'_3\rangle\langle S I_3, \mathbf{II_3} \mid S' I'_3\rangle, \tag{D2}$$

where $\mathbf{O}$ is an $SU_S(2)\times SU_I(2)$ irreducible tensor operator, and $\langle S' \mid\mid \mathbf{O}^{\mathbf{J}}_{\mathbf{I}} \mid\mid S\rangle$ is the reduced matrix element. Note that the notation $\mid\mid S\rangle$ indicates the spin-flavor states in the symmetric representation $(N_c,0,0)$ with $I=S$. The reduced matrix elements of the $SU(4)$ generators read:

$$\begin{aligned}
\langle S' \mid\mid S \mid\mid S\rangle &= \delta_{SS'}(2S+1)\sqrt{S(S+1)},\\
\langle S' \mid\mid I \mid\mid S\rangle &= \delta_{SS'}(2S+1)\sqrt{S(S+1)},\\
\langle S' \mid\mid G \mid\mid S\rangle &= \delta_{\{S,S',1\}}\frac{1}{4}\sqrt{(2S+1)(2S'+1)}\zeta(N_c,S,S'),
\end{aligned} \tag{D3}$$

where $\delta_{\{S,S',1\}}=1$ if $\mid S-S'\mid\leq 1$ and otherwise vanishes.

Reduced matrix elements of the operators involving the projects $\mathcal{P}_n$ are easily obtained using that $\mathcal{P}_n=\sum_{S_{n3},I_{n3}}\mid S_n, S_{n3}I_{n3}\rangle\langle S_n, S_{n3}I_{n3}\mid$, and the $SU(2)$ re-coupling results [70]. For the masses the relevant such RME becomes:

$$\langle S \mid\mid G^{ia}\mathcal{P}_n G^{ia} \mid\mid S\rangle = \frac{1}{2S+1}\langle S_n \mid\mid G \mid\mid S\rangle^2. \tag{D4}$$

For the axial currents the following RME is needed, namely:

$$\begin{aligned}
\langle S' \mid\mid G^{jb}\mathcal{P}_{n'}G^{ia}\mathcal{P}_n G^{jb} \mid\mid S\rangle \;&=\; \begin{Bmatrix} S & S_n & 1\\ S_{n'} & S' & 1\end{Bmatrix}^2\\
&\times\; \langle S' \mid\mid G \mid\mid S_{n'}\rangle\langle S_{n'} \mid\mid G \mid\mid S_n\rangle\langle S_n \mid\mid G \mid\mid S\rangle.
\end{aligned} \tag{D5}$$

Various reduced matrix elements which appear in the evaluation of the UV divergent pieces of the one-loop contributions to the self energy and to the axial currents are given below. They are obtained using the results given in the above Eqs. (D4) and (D5):

$$\langle S' \mid\mid G^2 \mid\mid S\rangle = \frac{1}{2}\delta_{SS'}(2S+1)(-S(S+1)+\frac{3}{8}N_c(4+N_c)), \tag{D6}$$

where $G^2 = G^{ia}G^{ia}$.

$$\langle S' \mid\mid G\hat{S}^2G \mid\mid S\rangle = \frac{1}{2}\delta_{SS'}(2S{+}1)\left(\frac{3}{4}N_c(4+N_c) + \frac{1}{8}(-40+12N_c+3N_c^2)S(S+1) - (S(S+1))^2\right), \tag{D7}$$

$$\begin{aligned}\langle S' \mid\mid G\hat{S}^4G \mid\mid S\rangle \ &= \ \frac{1}{2}\delta_{SS'}(2S+1)\left(\frac{3}{4}N_c(4+N_c) + (-6+5N_c+\frac{5}{4}N_c^2)S(S+1)\right.\\ &+ \ \left.(-7+\frac{3}{4}N_c+\frac{3}{16}N_c^2)(S(S+1))^2 - \frac{1}{2}(S(S+1))^3\right).\end{aligned} \tag{D8}$$

In the following, $S' = S$ or $S \pm 1$. With obvious notation:

$$\langle S' \mid\mid \{S^i, G^{ia}\} \mid\mid S\rangle = \delta_{SS'}(1+\frac{N_c}{2})(2S+1)\sqrt{S(S+1)}, \tag{D9}$$

obtained using the corresponding reduction relation in Table II.

$$\langle S' \mid\mid S^iI^a \mid\mid S\rangle = \delta_{SS'}(2S+1)S(S+1), \tag{D10}$$

$$\langle S' \mid\mid SG^{ia}S \mid\mid S\rangle = \frac{1}{2}(S(S+1)+S'(S'+1)-2)\langle S' \mid\mid G \mid\mid S\rangle, \tag{D11}$$

$$\begin{aligned}\langle S' \mid\mid GG^{ia}G \mid\mid S\rangle \ &= \ \frac{1}{16}\langle S' \mid\mid G \mid\mid S\rangle\\ &\times \ (3N_c(4+N_c) - 4(2+S(S+1)+S'(S'+1))), \end{aligned} \tag{D12}$$

$$\begin{aligned}\langle S' \mid\mid GG^{ia}\hat{S}^2G \mid\mid S\rangle \ &= \ \frac{1}{2}\langle S' \mid\mid G \mid\mid S\rangle\left(\frac{3}{4}N_c(4+N_c)\right.\\ &+ \ S(S+1)(-S(S+1)-5+\frac{3}{8}N_c(4+N_c)+\frac{1}{2}S'(S'+1))\\ &- \ \left.S'(S'+1)(1+\frac{1}{2}S'(S'+1)\right), \end{aligned} \tag{D13}$$

$$\begin{aligned}\langle S' \mid\mid GG^{ia}\hat{S}^4G \mid\mid S\rangle \ &= \ \frac{1}{2}\langle S' \mid\mid G \mid\mid S\rangle\\ &\times \ \left(\frac{3}{2}N_c(4+N_c) + (S(S+1))^2(-16+\frac{3}{8}N_c(4+N_c)-3S'(S'+1))\right.\\ &+ \ (S'(S'+1))^2(1-\frac{3}{2}S'(S'+1))\\ &+ \ \left.S(S+1)(-12+\frac{5}{2}N_c(4+N_c)+\frac{7}{2}(S'(S'+1))^2)\right), \end{aligned} \tag{D14}$$

$$\begin{aligned}\langle S' \mid\mid G\hat{S}^2G^{ia}\hat{S}^2G \mid\mid S\rangle \ &= \ \frac{1}{4}\langle S' \mid\mid G \mid\mid S\rangle\\ &\times \ \left((2+N_c)^2 + \frac{1}{2}(-16+5N_c(4+N_c))(S(S+1)+S'(S'+1))\right.\\ &- \ (S(S+1))^2(9+S(S+1)) - (S'(S'+1))^2(9+S'(S'+1))\\ &+ \ \left.\frac{3}{4}(-16+N_c(4+N_c))S(S+1)S'(S'+1)\right). \end{aligned} \tag{D15}$$

Finally, using that for any spin and isospin singlet operator (not necessarily an $SU(4)$ singlet) $\mathcal{Q}$, $\langle S' \mid\mid \mathcal{Q}\mathbf{O}^{\mathbf{J}}_{\mathbf{I}} \mid\mid S\rangle = \frac{1}{2S'+1}\langle S' \mid\mid \mathcal{Q} \mid\mid S'\rangle\langle S' \mid\mid \mathbf{O}^{\mathbf{J}}_{\mathbf{I}} \mid\mid S\rangle$, one can easily obtain the rest of the matrix elements involved in the calculation of the axial currents.

[1] H. Pagels, Phys.Rept. **16**, 219 (1975).

[2] S. Weinberg, Phys.Rev. **166**, 1568 (1968).

[3] S. R. Coleman, J. Wess, and B. Zumino, Phys.Rev. **177**, 2239 (1969).

[4] J. Callan, Curtis G., S. R. Coleman, J. Wess, and B. Zumino, Phys.Rev. **177**, 2247 (1969).

[5] J. Gasser, M. Sainio, and A. Svarc, Nucl.Phys. **B307**, 779 (1988).

[6] V. Bernard, N. Kaiser, and U.-G. Meissner, Int.J.Mod.Phys. **E4**, 193 (1995), hep-ph/9501384.

[7] E. E. Jenkins and A. V. Manohar, Phys.Lett. **B255**, 558 (1991).

[8] P. J. Ellis and H.-B. Tang, Phys.Rev. **C57**, 3356 (1998), hep-ph/9709354.

[9] T. Becher and H. Leutwyler, Eur.Phys.J. **C9**, 643 (1999), hep-ph/9901384.

[10] T. Fuchs, J. Gegelia, G. Japaridze, and S. Scherer, Phys.Rev. **D68**, 056005 (2003), hep-ph/0302117.

[11] E. E. Jenkins and A. V. Manohar, Phys.Lett. **B259**, 353 (1991).

[12] T. R. Hemmert, B. R. Holstein, and J. Kambor, Phys.Lett. **B395**, 89 (1997), hep-ph/9606456.

[13] T. R. Hemmert, B. R. Holstein, and J. Kambor, J.Phys.G **G24**, 1831 (1998), hep-ph/9712496.

[14] T. R. Hemmert, M. Procura, and W. Weise, Phys.Rev. **D68**, 075009 (2003), hep-lat/0303002.

[15] N. Fettes and U. G. Meissner, Nucl.Phys. **A679**, 629 (2001), hep-ph/0006299.

[16] M. Procura, B. Musch, T. Wollenweber, T. Hemmert, and W. Weise, Phys.Rev. **D73**, 114510 (2006), hep-lat/0603001.

[17] C. Hacker, N. Wies, J. Gegelia, and S. Scherer, Phys.Rev. **C72**, 055203 (2005), hep-ph/0505043.

[18] V. Bernard, T. R. Hemmert, and U.-G. Meissner, Phys.Lett. **B565**, 137 (2003), hep-ph/0303198.

[19] V. Bernard, T. R. Hemmert, and U.-G. Meissner, Phys.Lett. **B622**, 141 (2005), hep-lat/0503022.

[20] M. Procura, B. Musch, T. Hemmert, and W. Weise, Phys.Rev. **D75**, 014503 (2007), hep-lat/0610105.

[21] G. 't Hooft, Nucl.Phys. **B72**, 461 (1974).

[22] E. Witten, Nucl.Phys. **B160**, 57 (1979).

[23] J.-L. Gervais and B. Sakita, Phys.Rev.Lett. **52**, 87 (1984).

[24] J.-L. Gervais and B. Sakita, Phys.Rev. **D30**, 1795 (1984).

[25] R. F. Dashen and A. V. Manohar, Phys.Lett. **B315**, 425 (1993), hep-ph/9307241.

[26] R. F. Dashen and A. V. Manohar, Phys.Lett. **B315**, 438 (1993), hep-ph/9307242.

[27] E. E. Jenkins, Phys.Rev. **D53**, 2625 (1996), hep-ph/9509433.

[28] R. Flores-Mendieta, E. E. Jenkins, and A. V. Manohar, Phys.Rev. **D58**, 094028 (1998), hep-ph/9805416.

[29] R. Flores-Mendieta and C. P. Hofmann, Phys.Rev. **D74**, 094001 (2006), hep-ph/0609120.

[30] T. D. Cohen and W. Broniowski, Phys.Lett. **B292**, 5 (1992), hep-ph/9208253.

[31] P. Hagler, Phys.Rept. **490**, 49 (2010), 0912.5483.

[32] Z. Fodor and C. Hoelbling, Rev.Mod.Phys. **84**, 449 (2012), 1203.4789.

[33] C. Alexandrou, Prog.Part.Nucl.Phys. **67**, 101 (2012), 1111.5960.

[34] S. Durr, Z. Fodor, J. Frison, C. Hoelbling, R. Hoffmann, et al., Science **322**, 1224 (2008), 0906.3599.

[35] A. Walker-Loud, H.-W. Lin, D. Richards, R. Edwards, M. Engelhardt, et al., Phys.Rev. **D79**, 054502 (2009), 0806.4549.

[36] S. Aoki et al. (PACS-CS Collaboration), Phys.Rev. **D79**, 034503 (2009), 0807.1661.

[37] S. Aoki et al. (PACS-CS Collaboration), Phys.Rev. **D81**, 074503 (2010), 0911.2561.

[38] Y. Aoki et al. (RBC Collaboration, UKQCD Collaboration), Phys.Rev. **D83**, 074508 (2011), 1011.0892.

[39] W. Bietenholz, V. Bornyakov, M. Gockeler, R. Horsley, W. Lockhart, et al., Phys.Rev. **D84**, 054509 (2011), 1102.5300.

[40] R. Edwards et al. (LHPC Collaboration), Phys.Rev.Lett. **96**, 052001 (2006), hep-lat/0510062.

[41] J. Bratt et al. (LHPC Collaboration), Phys.Rev. **D82**, 094502 (2010), 1001.3620.

[42] C. Alexandrou et al. (ETM Collaboration), Phys.Rev. **D83**, 045010 (2011), 1012.0857.

[43] T. Yamazaki et al. (RBC+UKQCD Collaboration), Phys.Rev.Lett. **100**, 171602 (2008), 0801.4016.

[44] T. Yamazaki, Y. Aoki, T. Blum, H.-W. Lin, S. Ohta, et al., Phys.Rev. **D79**, 114505 (2009), 0904.2039.

[45] H.-W. Lin, T. Blum, S. Ohta, S. Sasaki, and T. Yamazaki, Phys.Rev. **D78**, 014505 (2008), 0802.0863.

[46] E. E. Jenkins and A. V. Manohar (1991).

[47] T. D. Cohen, Rev.Mod.Phys. **68**, 599 (1996).

[48] S. Weinberg, Nucl.Phys. **B363**, 3 (1991).

[49] J. Beringer et al. (Particle Data Group), Phys.Rev. **D86**, 010001 (2012).

[50] S. R. Beane and M. J. Savage, Phys.Rev. **D70**, 074029 (2004), hep-ph/0404131.

[51] S. R. Beane, Phys.Rev. **D70**, 034507 (2004), hep-lat/0403015.

[52] G. Colangelo, A. Fuhrer, and C. Haefeli, Nucl.Phys.Proc.Suppl. **153**, 41 (2006), hep-lat/0512002.

[53] A. A. Khan, M. Gockeler, P. Hagler, T. Hemmert, R. Horsley, et al., Phys.Rev. **D74**, 094508 (2006), hep-lat/0603028.

[54] P. F. Bedaque, H. W. Griesshammer, and G. Rupak, Phys.Rev. **D71**, 054015 (2005), hep-lat/0407009.

[55] W. Detmold and M. J. Savage, Phys.Lett. **B599**, 32 (2004), hep-lat/0407008.

[56] B. Smigielski and J. Wasem, Phys.Rev. **D76**, 074503 (2007), 0706.3731.

[57] K.-I. Ishikawa et al. (PACS-CS Collaboration), Phys.Rev. **D80**, 054502 (2009), 0905.0962.

[58] S. Beane, E. Chang, W. Detmold, H. Lin, T. Luu, et al., Phys.Rev. **D84**, 014507 (2011), 1104.4101.

[59] L.-s. Geng, X.-l. Ren, J. Martin-Camalich, and W. Weise, Phys.Rev. **D84**, 074024 (2011), 1108.2231.

[60] A. Calle Cordón (work in preparation).

[61] J. Green, M. Engelhardt, S. Krieg, J. Negele, A. Pochinsky, et al. (2012), 1209.1687.

[62] M. Procura, T. R. Hemmert, and W. Weise, Phys.Rev. **D69**, 034505 (2004), hep-lat/0309020.

[63] V. Bernard, Prog.Part.Nucl.Phys. **60**, 82 (2008), 0706.0312.

[64] L. Geng, J. Martin Camalich, L. Alvarez-Ruso, and M. Vicente Vacas, Phys.Rev. **D78**, 014011 (2008), 0801.4495.

[65] A. Calle Cordón and J. L. Goity (work in preparation).

[66] T. DeGrand, Phys.Rev. **D86**, 034508 (2012), 1205.0235.

[67] S. Scherer, Adv.Nucl.Phys. **27**, 277 (2003), hep-ph/0210398.

[68] C. E. Carlson, C. D. Carone, J. L. Goity, and R. F. Lebed, Phys.Rev. **D59**, 114008 (1999),

hep-ph/9812440.

[69] R. F. Dashen, E. E. Jenkins, and A. V. Manohar, Phys.Rev. **D51**, 3697 (1995), hep-ph/9411234.

[70] A. Edmonds, *Angular momentum in quantum mechanics*, Oxford Univ. Press, London (1955).